RECOMMENDED PREVENTIVE SEXUAL HEALTH SERVICES FOR PEOPLE WITH A VAGINA/VULVA

Sexual Health Service	Adolescent (13 to 18 years)	Adult (19 and older)
Cervical cancer screening	Not recommended	Those aged 21 to 29 every 3 years Those aged 30 to 65 every 3 to 5 years
		Not recommended for those aged 66 and over
Chlamydia screening	Those who are sexually active (screen annually) or pregnant	Those age 24 and under (screen annually)
		Those age 25 and over who are at risk*
		Those who are pregnant, if 24 and under, or pregnant and at risk if aged 25 and over
Contraceptive counseling	Those who are sexually active	Those who are sexually active and/or of childbearing age
Counseling to prevent sexually transmitted infections	Those who are sexually active	Those who are at risk*
Gonorrhea screening	Those who are sexually active or pregnant	Those aged 24 and under (screen annually), including those who are pregnant
		Those aged 25 and over who are at risk, including those who are pregnant
Hepatitis A vaccine	Those not vaccinated as children	Those who are at risk* and are unvaccinated
Hepatitis B vaccine	Those not vaccinated as children	Those who are at risk* and are unvaccinated
		Those born in a country where the hepatitis B virus is common
Hepatitis B screening	Those who are pregnant	Those who are pregnant
	Those who are at risk and are unvaccinated	Those who are at risk* and are unvaccinated
	Those born in a country where the hepatitis B virus is common	Those born in a country where the hepatitis B virus is common
Hepatitis C screening	Those who are age 18	Those aged 19 to 79
	Those who are pregnant	Those who are at risk* or pregnant
Herpes	Those who have been exposed or are symptomatic	Those who have been exposed or are symptomatic
HIV testing	Those aged 13 and older (test at least once) Those who are pregnant	Those who are aged 19 to 64 (test at least once)
		Those who are at risk* (test at least annually)
		Those who are pregnant
HPV vaccine	Those not vaccinated at age 11 or 12	Those not vaccinated through age 26 (and eligible adults aged 27 to 45)
Syphilis screening	Those who are pregnant or at risk	Those who are pregnant or at risk*
Trichomoniasis	Not recommended	Those who are at risk or receiving care in high-risk settings (correctional facilities, STI clinics)

^{*}You're at risk if you have had unprotected vaginal, anal, or oral sex (sex without a condom), have multiple partners, have an STI (including HIV), share injection drug equipment (including needles or syringes), exchange sex for drugs or money, are incarcerated, or have a partner who answers "yes" to any of these questions or whose health status you don't know.